



Rabbits

All employees and students working with animals should be aware that laboratory animals may bite or scratch. In addition, handling of cages, pens and other equipment could cause cuts or scratches. Cuts, bites and scratches could potentially expose the employees to viral, bacterial, parasitic or allergic agents, which are transmissible from animals to humans. Although the risk of bites and scratches is species dependent, there are a few simple guidelines, which if followed, will significantly reduce the potential risks of such incidents. These include:

- Always wear appropriate personal protective equipment, especially hand and face/eye protection.
- If moving large contaminated items (e.g. non-human primate cages), wear heavy gloves.
- When available and appropriate, use mechanical restrainers according to IACUC guidelines when performing procedures on unanesthetized animals.
- All bites and scratches that result in bleeding should be immediately and thoroughly scrubbed and cleansed with soap and running water for at least 15 minutes. Notify supervisor and seek medical attention immediately.
- There are some potential hazards inherent in any work environment. These include poor ergonomics, slips and falls, electrical safety hazards, etc. UTSouthwestern Medical Center has developed a wide range of environmental health and safety guidelines to address these potential hazards.

ALLERGY HAZARDS

One of the most common health concerns in the laboratory animal setting is a work-associated allergy. The risk of developing an allergy depends on parameters such as species, facility, ventilation, and the employee's "base-line" health status. There are also several individual risks which can be divided into four overlapping Risk Groups:

- **Normal:** No evidence of allergic disease;
- **Atopic:** Pre-existing allergic disease;
- **Asymptomatic:** Antibodies to animal allergens;
- **Symptomatic:** Clinical symptoms on exposure to allergenic animal proteins.
 - Symptoms of allergic reaction vary depending on the severity of the reaction and include:
 - **Contact urticaria** with symptoms such as: redness, itchiness of skin, welts and hives;
 - **Allergic conjunctivitis** with symptoms such as: itchy, burning, running, or red eyes.
 - **Allergic rhinitis** with symptoms such as: sneezing, itchiness, clear nasal drainage, nasal congestion;

UT SOUTHWESTERN MEDICAL CENTER
OCCUPATIONAL HEALTH DEPARTMENT
WORKER PROTECTION PROGRAM

- **Asthma** with symptoms such as: cough, wheezing, chest tightness, shortness of breath;
- **Anaphylaxis** with symptoms such as: generalized itching, hives, throat tightness, eye or lip swelling, difficulty in swallowing, hoarseness, shortness of breath, dizziness, fainting, nausea, vomiting, abdominal cramps, and diarrhea.

All Rodent Areas:

Conventional and barrier rodent environments have allergens present, such as dust from bedding and animal dander. High risk exposure activities include cage wash, cage changing, and experimental work with animals that haven't been anesthetized. Exposures are also higher in rooms without microisolator lids.

HOW TO PROTECT YOURSELF

- Wearing appropriate PPE and respiratory protection can help to limit the amount of animal dander exposure.

INFECTIOUS DISEASE HAZARDS

Laboratory rodents rarely transmit zoonotic infections anymore, because they are supplied by commercial vendors who eliminate these infections from their stock.

- ***Pasturella multocida***: *Pasteurella multocida* is a serious pathogen of rabbits and has been associated with infected bites and scratches sustained from rabbits.
 - **Reservoir/source of infection**: Many mammals, including rabbits
 - **Transmission**: Bite wounds
 - **Disease in people**: Cellulitis at site of bite and sepsis may occur
- ***Salmonellosis***
 - **Reservoir/source of infection**: Many mammals, including rabbits.
 - **Transmission**: Fecal/Oral
 - **Disease in people**: Gastroenteritis with diarrhea
- ***Tularemia*** - The bacteria that causes tularemia can infect wild rabbits, but is not found in laboratory rabbits obtained through commercial vendors.
- ***Other infections***: Skin fungi and mites carried by rabbits can be transmitted to humans causing a rash.

HOW TO PROTECT YOURSELF

- Until you have washed your hands, keep them away from your mouth, nose and eyes.
- Do not eat, drink or store food in research areas.
- Do not handle contact lenses in the laboratory. Wearing of gloves, lab coat or scrubs is required. Other personal protective equipment such as dust/mist masks or approved respirator masks (e.g., Type N95 by 3M company) are strongly recommended when working with animals.

PHYSICAL HAZARDS

- ***Bites and scratches***: The potential for receiving a bite or scratch is an ever-present hazard that faces all employees working directly with laboratory animals and related

Craig Glazer MD
Medical Director
Worker Protection Program

Tam Johnson RN, BSN, COHN-S
Nurse Manager
Occupational Health

UT SOUTHWESTERN MEDICAL CENTER
OCCUPATIONAL HEALTH DEPARTMENT
WORKER PROTECTION PROGRAM

equipment. Employees should be properly trained in handling and general restraint techniques of the species they are assigned to. Additionally, all staff should be familiar with first aid procedures specific to each species

- *Needlestick / Sharps injury*

HOW TO PROTECT YOURSELF

- Always wear appropriate personal protective equipment, especially hand and face/eye protection.
- If moving large contaminated items, wear heavy gloves.
- Proper training and awareness of animal behavior are vitally important to avoid injuries.
- When available and appropriate, use mechanical restrainers when performing procedures on unanesthetized animals.
- All bites and scratches that result in bleeding should be immediately and thoroughly scrubbed and cleansed with soap and running water for at least 15 minutes.
- Read and understand the protocol-related procedures before you start the experiment. If necessary, do a dry run.
- Do not recap needles; dispose of them in appropriate sharp containers.
- Use safety needles and sharp devices and avoid sharps whenever possible by substituting manually operated pipettes for needles and syringes, and cannulae for needles.
- **Until you have washed your hands, keep them away from your mouth, nose and eyes.**
- Read and understand the special safety requirements for each work area or animal species.
- Clean all spills immediately.
- Report all incidents or equipment malfunctions to your supervisor immediately

PROTOCOL RELATED HAZARDS

Protocol-related hazards are defined as those specifically associated with either routine operational or experiment-specific protocols. Some general hazards also associated with protocols, such as the risk of fire in the use of bunsen burners or torches, or electrical hazards in the use of experiment-specific equipment, are not included in the category of protocol-related hazards. Protocol-related hazards are protocol-specific. For example:

- Hazards associated with the use of a specific viral vector carrying a transgene for toxin production or with a piece of prototype equipment to perform an experimental task.
- Experimental studies can involve any number of hazards such as the use of radioactive materials, infectious agents, toxins or toxic chemicals, flammable substances, etc.

HOW TO PROTECT YOURSELF

- Follow the steps in your approved safety plan.
- Educate yourself on the protocol specific hazards.

Craig Glazer MD
Medical Director
Worker Protection Program

Tam Johnson RN, BSN, COHN-S
Nurse Manager
Occupational Health

UT SOUTHWESTERN MEDICAL CENTER
OCCUPATIONAL HEALTH DEPARTMENT
WORKER PROTECTION PROGRAM

Additional information regarding various protocol-related hazards may be obtained from

www.utsouthwestern.edu/workerprotection

Craig Glazer MD
Medical Director
Worker Protection Program

Tam Johnson RN, BSN, COHN-S
Nurse Manager
Occupational Health